ORIGINAL PAPER



Everything in Moderation: Moderate Use of Screens Unassociated with Child Behavior Problems

Christopher J. Ferguson¹

Published online: 7 February 2017 © Springer Science+Business Media New York 2017

Abstract The impact of children's use of "screen" media including television and computer games, continues to be debated. The American Academy of Pediatrics (AAP) until recently recommended a relatively restrictive screen time diet of 2 h or less for most youth. A representative correlational sample of youth were assessed for links between screen time and risky behavioral outcomes. Data collection occurred in 2013 conducted by the State of Florida. Use of screens that was moderately high, in excess of the AAP's former recommendations, but not excessive (1 SD or higher than average), was not associated with delinquency, risky behaviors, sexual behaviors, substance abuse, reduced grades or mental health problems. Even excessive screen use (1 SD or higher) was only weakly associated with negative outcomes related to delinquency, grades and depression only, and at levels unlikely to be practically significant. Results conceptually replicate those of Przybylski (2014) with a US sample for depression and delinquency as outcomes. Moderate use of screens, though in excess of the AAP's historical recommendations, are unassociated with problem outcomes. Excessive use of screens is only weakly associated with negative outcomes, and only those related to depression and delinquency as well as reduced grades, but not risky driving, substance use, risky sex or disordered eating. Although an "everything in moderation" message when discussing screen time with parents may be most productive, results do not support a strong focus on screen time as a preventative measure for youth problem behaviors.

Keywords Video games · Television · Screen time · Adolescents · Risk behaviors

Introduction

Screen technology, ranging from televisions to computers to smart phones have become increasingly integrated into the lives of youth (minors age 12–18). Considering all screens

Christopher J. Ferguson CJFerguson1111@aol.com

¹ Department of Psychology, Stetson University, 421 N. Woodland Blvd, DeLand, FL 32729, USA

together, it is estimated that screen time among youth averages approximately 7–8 h per day [1]. Organizations such as the American Academy of Pediatrics typically advocate for far fewer screen hours for youth, typically around 2 h a day. However, with screens becoming increasingly integrated into education, work, communication and socialization, rather than just passive entertainment, the applicability of such guidelines has become less clear. For instance, with younger children, the organization Zero-to-Three has noted that how screens are used is often more critical than simply focusing on number of hours [2]. Nonetheless, concerns persist regarding the potential impact of excessive screen time (defined here as 1 SD above mean use) on youth behavioral outcomes.

Relationships between screen time and behavioral outcomes have not always been clear. Although most scholars would likely agree that screen time should be balanced with other activities and excessive use avoided, disagreement exists on the importance of screen time as a predictor of behavioral issues. Evidence on outcomes is often mixed, with suggestions of complexity in the interaction between screens and behavior. For example, screen time is associated with obesity, but only to the extent it interferes with physical activity. If recommendations for physical activity are met, screen time, even if considerable, no longer predicts obesity [3]. Likewise, evidence suggests that links between screen time and cognitive performance are complex, often relating to quality more than quantity [4], without clear evidence that screen time exposure relates to test scores [5].

Regarding behavioral issues, research has had difficulty observing linear consistent linear relationships between screen time and outcomes such as criminal aggression [6], disordered eating [7], and depression [8]. One possibility is that screen time influences may not be linear at all but may instead be dose dependent. That is to say that screen time consumption that is within the average range may be developmentally normal, with only excessive use compared to peers being indicative of potential problems. This approach differs from the historically typical screen time recommendations with absolute time cut-offs, above which use may be considered to be problematic no matter how normative.

Several recent studies have indicated support for the dose-dependent explanation for links between screen time and problem outcomes. Przybylski [9] examined the relationship between time spent playing video games and externalizing and internalizing symptoms in a large sample of children from the UK. Light users of video games were found to have the fewest symptoms, with non-users and moderate users (those with 1 SD of mean use) also relatively low in symptoms. Heavy use of video games was associated with more externalizing and internalizing symptoms, although Przybylski observed that, even for heavy users, the relationship between video game time and behavioral outcomes was very small, potentially lacking practical significance. Such dose dependent effects have been found in other studies, often indicating worse results for both heavy users and non-users of video games [10, 11].

Few studies have examined this issue for screen time more generally nor with a wider array of behavioral outcomes. As such this study will examine dose-dependent influences of screen time on a range of behavioral outcomes including depressive symptoms, delinquency, risky driving, substance abuse, risky sex, disordered eating and GPA (Grade Point Aveage). It is hypothesized in each case that excessive (1 SD above the mean), but not moderate (within 1 SD of the mean) or light screen time will be associated with negative behavioral outcomes despite other relevant factors being controlled.

Methods

Participants

Participants in the current study were a representative sample of 6089 youth from the state of Florida. Participants were part of the 2013 Youth Risk Behavior Survey which is coordinated by the Centers for Disease Control. Participants were almost equally distributed in terms of sex (50.9% were female). Regarding race, the largest percentage of youth were Caucasian (43.7%) with smaller amounts of African-American (22.8%), Native American (0.4%), Hispanic (27.7%) Asian (2.0%) or other groups (3.4%). The average age of our sample was 16 (range 12–18, SD = 1.24).

Materials

Data in the current study were drawn from the Florida version of the Youth Risk Behavior Surveillance System's school-based Youth Risk Behavior Survey (YRBS). Description of the development and validation of the YRBS can be found from the CDC [12]. This survey instrument included several scales of interest to the present study. In many cases scale metrics differed between questions. As such all answers were first converted to z-scores prior to summation into scale scores. Reported coefficient alphas are for the current sample. These included the following:

Risk/Protective Factor Variables

Sleep A single item "on an average school night, how many hours of sleep do your get?" was included to measure sleep quality of youth. This variable can indicate overall stress, and quality of rest which can influence mental health.

Physical Activity Three items measuring overall physical activity, participation in PE (physical education) and participation in sports (alpha = .53) were included as a measure of physical activity. Physical activity for youth is normative and developmentally healthy and time spent on physical activity takes time away from risky behaviors.

Family Involvement Family involvement was measured by a single item inquiring as to the frequency of the youth's family eating dinner together. This was included as a general indicator of family cohesiveness.

Depression This scale consisted of three items related to depressive symptoms. "During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?" is an example item. Coefficient alpha was .72. Depression is used both as an outcome on its own, as well a risk factor for other outcomes, as previous research has indicated that risk taking behaviors are more common with children experiencing depression [13].

Screen Time Two items measured frequency of television and video game screen time. "On an average school day, how many hours do you watch TV" is an example item. These were summed together to form a measure of total media viewing time.

Outcome Variables

Depression As noted above, depression was included as an outcome variable in its own right as well as a risk factor for other outcomes.

Delinquency Four items were used to measure delinquency. These concerned themselves with issues of weapons carrying, fighting off and on school property, and being injured in a physical fight. Alpha reliability was .76

Risky Driving Three items concerned themselves with risky driving behaviors (alpha = .68). These items involved driving and drinking, riding in a car while the driver had been drinking and texting while driving. A fourth item planned for inclusion in the preregistration (as noted below, the study was preregistered) regarding seat belt use was dropped due to low reliability with the other items.

Illegal Substance Use This eleven item scale measured the frequency of use of multiple forms of illegal drugs (from marijuana to "hard" drugs such as cocaine and heroin). Coefficient alpha was .91.

Risky Sexual Behaviors This five item scale (alpha = .89) examined risky sexual behaviors the youth engaged in. These included lifetime and recent sexual partners, condom use, sex while intoxicated and pregnancy.

Disordered Eating Disordered eating was assessed via a three item scale (alpha = .58). Items involved purposeful weight loss using fasting, pills or vomiting. An initial plan to include healthy eating as an outcome in the preregistration was not included, to maintain focus on risky behaviors. Note that this outcome was not assessed prior to making this decision.

Grades (GPA) Students were asked to provide their grades using a single item.

Procedure

Participation in the current study was achieved through two-step cluster probability sampling. First, schools in Florida were randomly selected for participation, then classroom within those schools were randomly selected for participation. All students in the selective classrooms were invited to participate. This procedure was designed to produce a sample representative of youth attending schools in the state of Florida. Passive parental consent was used for all youth in grades 9 through 12, and children assented to the procedure. All procedures were designed to meet federal standards for ethical research with human participants. Although the data is archival a data and analysis plan was preregistered prior to analysis and is available at: https://osf.io/7xm3j/

Results

All results were analyzed using OLS (ordinary least squares) regression using SPSS software. Predictor variables were entered first into the equation, followed by screen time

contrasts. Consistent with Przybylski [9] screen time contrasts were developed to examine for potential dosing effects. These consisted of four groups. First were abstainers, those youth who reported no screen time whatsoever. Low users were those youth who fit more closely with the standard guidelines of no more than 2 h a day of screen time. Moderate users included those users within a standard deviation of the mean on screen time. Moderate users would correspond to roughly three to six hours of use each day. Excessive users included all users at least one standard deviation above the mean on screen time. Contrasts were dummy coded, with abstainers used as the reference category. Collinearity diagnostics revealed an absence of issues related to multicollinearity in the current analysis, with VIFs typically no higher than 1.12.

All results are presented in Table 1. Across most outcomes male gender and depression were significant risk factors for worse outcomes, consistent with prior research. Family involvement was a fairly consistent protective factor.

Regarding screen time, neither low to moderate use of screens was related to worse outcomes than was media abstinence. This corresponds to the observation that up to six hours of screen time as indicated by TV and computer game use, was not a risk factor for negative outcomes. Excessive use was significantly related to depression, delinquency and reduced grades, but not to other outcomes. Thus, results of Przybylski [9] are replicated here regarding depression and delinquency, but not extended to other risky outcomes aside from grades. However, also consistent with Przybylski, the effect sizes for outcomes related to screen time were very small, typically accounting for less than half a percent of variance in the outcome variable (in the case of delinquency), to just over 1% variance overlap in the case of depression. In the event of such small effect sizes, the risk of overinterpretation of outcomes is significant. Thus, consistent with the approach recommended by Przybylski, outcomes in this effect size range, even for excessive screen time, suggest that screen time is not an efficient risk factor for negative outcomes.

Discussion

Screen time among children continues to be a hotly debated issue, with conflicting views about the proper recommendations to give to children regarding screen time. At present,

Control Variables	Depression	Delinquency	Risky Driving	Substance. Use	Risky Sex	Eating	GPA
Male Gender Sleep Physical Activity Family Involvement	14*** 16*** 01 10***	.22*** 05 .05 07 15***	.08*** 05 .09*** 13***	.12*** 03 .04 11***	.12*** 08*** .07 16*** 12**	05 06 .01 13***	17*** .05 .03 .07
Media Contrasts Abst. v Low Abst. v Moderate Abst. v Excessive	02 .05 .13***	03 01 .07***	02 05 .02	01 01 .06	02 01 .06	05 04 03	.05 02 11***

Table 1 Associations between screen time, control variables and youth outcomes

Effect sizes reported for control variables are for the Abstinence versus Low use contrast. However, outcomes did not differ significant for other contrasts for control variables. All regression coefficients are standardized ***p < .001

a considerable gulf exists between the best known screen time recommendations provided by the American Academy of Pediatrics, and the actual screen use behaviors of youth. More crucially, screen time recommendations often remain best guesses, with data about the utility of such recommendations not always clear. This study sought to address gaps on previous literature by examining what levels of screen time were associated with negative outcomes and how strong these associations were. This data may help to guide organizations concerned with screen time recommendations to revise those recommendations in the future.

The present study conceptually replicated results of Przybylski [9] with depression and delinquency. Specifically, relations between screen time and negative outcomes were dose dependent, with exposures in considerable excess of the AAP's historical 2-h maximum recommendation required before associations with negative outcomes were noticeable. Specifically, negative outcomes were elevated only among youth who consumed over six hours of media a day, effectively outliers even among their peers. Further, even where these associations were noted, they were very small, accounting only for between 0.49% of the variance in delinquency and 1.7% of the variance in depressive symptoms. Thus it is unclear whether such small associations warrant the degree of attention they often receive from professional advocacy organizations. Or, put simply, a focus on screen time may not return as much "bang for the buck" as some may hope.

Unexpectedly, although results of the current study conceptually replicated those of Przybylski for delinquent and depressive, outcomes, they did not extend to most other outcomes. Among other outcomes, screen time only negatively predicted GPA, and once again the relationship was very small with screen time only predicting 1.2% of the variance in grades. For other outcomes related to risky driving, risky sex, substance abuse and restrictive eating, media was not a significant predictor with other life factors controlled.

By contrast gender was one of the most consistent predictors of negative outcomes. Male gender was generally predictive of most negative outcomes, with the exception of female gender which predicted depressive symptoms. Further, family involvement was generally successful as a protective predictor of negative outcomes. Family involvement in children's lives reduced their involvement in risky behaviors or other negative outcomes.

Policy Implications

Results from the current study have several potential policy implications. First, the current narrative on media effects among professional advocacy organizations may be too extreme. Arguably, most media related policy statements from the AAP and similar groups advocate for the potential for media, whether general use, or consumption of specific content, to have strong deleterious effects. At present, however, data to support such narratives is limited. This is not to say that professional advocacy organizations should not emphasize moderation. However, it may be inappropriate to needlessly frighten parents with the specter of outcomes related to violent crime, mental health problems, serious school failure, drug and alcohol abuse or eating disorder diagnoses, when evidence for substantial links between media use and these outcomes is fairly limited. Policy statements may be on firmer ground when emphasizing potential, though limited, impact on grades, obesity, or exclusion from other activities.

However, adopting a more moderate and measured tone may increase the credibility of such policy statements.

Second, it is likely the case that setting hard time limits on screen use is a fraught avenue for policy. First, data from the current study suggests that children are resilient to screen consumption at much higher levels, up to six hours daily, than is typically recommended by most policy statements. Thus, policy statements focused on restrictive media consumption may do more to foster guilt in parents unable to meet unrealistic expectations than they do help children. This may also apply to controversial areas of media use such as violent or sexy media, where policy statements often advocate abstinence or substantial restriction, a media use outcome that is unlikely to be met for most children, and for which data on efficacy are unclear at best. An unnecessary focus on objectionable content may ultimately devolve into moralizing rather than providing practically useful guidelines.

In addition, as noted by the group Zero to Three [2], focusing mainly on time consumption ignores the issue that *how* media are used is often more critical than how often media are used. Most existing media policy statements imply that media use is of limited value and, at best, detracts from engagement in other activities involving socialization or learning. However, this understanding of media may be unnecessarily simplistic as learning and socialization can often occur within the context of using media. Further, screens of various sorts are increasingly embedded into daily life, whether education, work, socialization or personal organization. Thus, narrow limits on screen time may not keep up with the myriad ways in which screens have become essential to modern life. Discouraging youth from becoming intimately familiar with screen technologies may ultimately do youth more disadvantage than advantage.

In general, professional advocacy organizations may wish to consider releasing fewer policy statements of all types. For instance, at present, the AAP has approximately eight active policy statements (despite such policy statements expire within 5 years) on various media, most of which focus on potential deleterious effects. This may inadvertently foster the impression of organizations such as the AAP as "anti-media" particularly when such policy statements do not disclose research that would conflict with their positions. It is unlikely that so many policy statements focused on deleterious outcomes are particularly informative for the general public or are in the long-term interest of groups like the AAP, to the extent they may alienate a general public disinclined to listen to media worriers. Given that scholars argue that it is unrealistic for pediatricians to impart the plethora of warnings and cautions advised by the AAP to parents within the limited time available for well-visits [14], the AAP may wish to consider retiring some of its media-based policy statements.

Limitations and Future Directions

As with most studies, the present one has limitations that reduce confidence in the ability to draw causal connections. First and foremost is the correlational nature of the current data which eliminate the potential for causal inferences. Secondly, the surveys were not designed with the specific current research questions in mind. As such, constructs are developed from preexisting pools of survey responses. Lastly, the current study draws data from only youth respondents. Having multiple respondents would be far more desirable.

By contrast, this study has, as a strength, the fact that it was preregistered, eliminating the potential that post-hoc data manipulation could result in spurious results. Given ongoing controversies about the methods in much of media effects research, increasingly adoption of

preregistered studies and other principles of open science could be illustrative in understanding what media effects are reliable, and which may be the product of historical cycles of societal concern over media. Further studies would also do well to focus on issues such as time spent on screens, or objectionable content, and more on how screen use becomes embedded in the social fabric of families and peer networks.

Concluding Statements

Debates about screen time are unlikely to abate in the near future. Current data suggests that screen time has limited impact on children's well-being. Counseling moderation in screen use is likely to result in the healthiest outcomes, while understanding that this should focus on peer normative screen use, rather than arbitrarily low cut-offs. Further, understanding the context of screen use and how it is embedded in the social fabric of modern life may be crucial in reenvisioning the role of screen use in young lives. Fundamentally, to the degree society is concerned about the well-being of youth, focusing on screen technology may be more distraction than panacea. This should be communicated more clearly, so that societal efforts can be refocused on issues that have more dramatic impact on young lives.

Compliance with Ethical Standards

Conflict of Interest The author has no conflicts of interest to declare and is solely responsible for the study design, analysis and decision to submit for publication. The author received no grants, honorarium or other funding to contribute to this report.

Human and Animal Rights All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in the study.

References

- Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. Sleep Medicine Reviews. June 2015;21:50–58.
- Zero to Three. Screen sense: Setting the record straight. Retrieved from: http://www.zerotothree. org/parenting-resources/screen-sense/
- Laurson K, Lee J, Eisenmann J. The cumulative impact of physical activity, sleep duration, and television time on adolescent obesity: 2011 youth risk behavior survey. *Journal of Physical Activity & Health*. March 2015;12(3):355–360.
- Linebarger D, Vaala S. Screen media and language development in infants and toddlers: an ecological perspective. *Developmental Review*. June 2010;30(2):176–202.
- Gentzkow M, Shapiro J. Preschool television viewing and adolescent test scores: historical evidence from the Coleman study. *The Quarterly Journal of Economics*. 2008; 123: 279–323.
- Schwartz J, Beaver K. Revisiting the association between television viewing in adolescence and contact with the criminal justice system in adulthood. *Journal of Interpersonal Violence*. In Press.
- 7. Holmstrom A. The effects of the media on body image: a meta-analysis. *Journal Of Broadcasting & Electronic Media.* 2004;48(2):196–217.
- Hittner J. How robust is the Werther effect? A re-examination of the suggestion-imitation model of suicide. Mortality: August 2005;10(3):193–200.
- 9. Pryzbylski A. Electronic gaming and psychosocial adjustment. Pediatrics. 2014

- Allahverdipour H, Bazargan M, Farhadinasab A, & Moeini B. Correlates of video games playing among adolescents in an Islamic country. *BMC Public Health*. 2010; 10: 286.
- Kutner L, Olson C. Grand theft childhood: the surprising truth about violent video games and what parents can do. New York: Simon & Schuster; 2008.
- Centers for Disease Control and Prevention. Methodology of the youth risk behavior surveillance system. MMWR 2004 53(No. RR-12):1–14.
- Anatale K, Kelly S. Factors influencing adolescent girls' sexual behavior: a secondary analysis of the 2011 youth risk behavior survey. *Issues In Mental Health Nursing*. 2015;36(3):217–221.
- Belamarich PF, Gandica R, Stein RE, Racine AD. Drowning in a sea of advice: pediatricians and American Academy of Pediatrics policy statements. *Pediatrics*. 2006;118(4). doi:10.1542/peds.2006-0652

Christopher J. Ferguson is a professor of psychology at Stetson University. He has conducted research on a wide range of media effects, including video game violence effects. He lives in Florida with his wife and son.

Psychiatric Quarterly is a copyright of Springer, 2017. All Rights Reserved.